

Colonoscopy Prep Instructions – 2-DAY MIRALAX GATORADE

Please follow the General Colonoscopy instructions along with your prep instructions.

Purchase the preparation materials:

- One (1) 238-gram bottle of Miralax Laxative (over-the-counter)
- One (1) box of 5 mg Dulcolax laxative tablets (20 mg needed) (NOT Dulcolax stool softener)
- Two (2) 32 oz (one quart) bottles of Gatorade
- If you are a diabetic, purchase G2 Reduced Sugar Gatorade (NOT red, purple, or blue)
You may also use water or Crystal Light Lemonade if you prefer.
- One (1) 10 oz bottle of Magnesium Citrate - CLEAR ONLY; keep in refrigerator

PM procedures can have breakfast 2 days prior to procedure.

Mix the entire 238-gram bottle of Miralax with 64oz of Gatorade or Crystal Light in a large pitcher. Be sure to stir the solution until the Miralax is completely dissolved and then place in the refrigerator. **DO NOT MIX WITH RED, PURPLE or BLUE Liquid.**

Begin 2 days before your procedure:

Drink clear liquids only all day. Try and drink 8oz of liquid every hour.

Take one 10oz bottle of Magnesium Citrate (clear only) at 6pm.

Day before your procedure:

Take 4 Dulcolax tablets (20mg) total at 4pm.

Begin at 6PM the day before your procedure:

Drink the first (32oz) of your mixture. Drink an 8oz glass every 10-15 minutes until the first 32oz have been completed. Refrigerate the remaining 32oz of mixture and continue to drink clear liquids to stay hydrated.

Begin 5 hours prior to procedure:

Drink the remaining 32oz of the mixture. Drink an 8oz glass every 10-15 minutes until the remaining mixture is gone.

All fluids must be stopped 2 hours prior to your arrival at our facility or your procedure may be cancelled.